

# Contents

<b>Acknowledgement</b>	<b>i</b>
<b>Abstract</b>	<b>ii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Quantum spin liquids . . . . .	1
1.2 Kitaev quantum spin liquid . . . . .	3
1.2.1 Kitaev model and the exact solution . . . . .	3
1.2.2 Fractional excitations and phase diagram . . . . .	5
1.2.3 Kitaev interactions in real materials . . . . .	6
1.2.4 Stability of the Kitaev spin liquid . . . . .	9
1.3 Candidate materials for the Kitaev spin liquid . . . . .	11
1.3.1 $\text{Na}_2\text{IrO}_3$ , $\alpha\text{-Li}_2\text{IrO}_3$ , and $\text{H}_3\text{LiIr}_2\text{O}_6$ . . . . .	11
1.3.2 $\alpha\text{-RuCl}_3$ . . . . .	12
1.3.3 $\beta$ - and $\gamma\text{-Li}_2\text{IrO}_3$ . . . . .	13
1.4 Signatures of the Kitaev quantum spin liquid . . . . .	14
1.4.1 Thermal fractionalization . . . . .	14
1.4.2 Dynamical spin structure factor in the ground state . . . . .	18
1.4.3 Raman scattering . . . . .	20
1.5 Purpose of this thesis . . . . .	22
1.6 Organization of this thesis . . . . .	24
<b>2 Model</b>	<b>25</b>
2.1 Jordan-Wigner transformation . . . . .	25
2.2 Majorana fermion representation . . . . .	27
<b>3 Method</b>	<b>29</b>
3.1 Quantum Monte Carlo method . . . . .	29
3.2 Cluster dynamical mean-field theory . . . . .	31
3.3 Continuous-time quantum Monte Carlo method . . . . .	34
3.4 Maximum entropy method . . . . .	37
3.5 Physical quantities . . . . .	38
<b>4 Results by CDMFT + CTQMC method</b>	<b>41</b>
4.1 Benchmark . . . . .	41
4.1.1 CDMFT method . . . . .	41

4.1.2	CDMFT+CTQMC method . . . . .	43
4.2	Dynamical spin structure factor . . . . .	45
4.3	NMR relaxation rate . . . . .	53
4.4	Magnetic susceptibility . . . . .	55
4.5	Korringa ratio . . . . .	56
4.6	Effect of the $Z_2$ variables $\{\eta_r\}$ on the dynamical properties . . . . .	60
4.7	Summary of this chapter . . . . .	63
<b>5</b>	<b>Results by QMC + CTQMC method</b>	<b>65</b>
5.1	Results for 2D honeycomb structure . . . . .	66
5.1.1	Magnetic susceptibility and NMR relaxation rate . . . . .	66
5.1.2	Dynamical spin structure factor . . . . .	68
5.2	Results for 3D hyperhoneycomb structure . . . . .	72
5.2.1	Magnetic susceptibility and NMR relaxation rate . . . . .	72
5.2.2	Dynamical spin structure factor . . . . .	73
5.3	Comparison between 2D and 3D results . . . . .	76
5.4	Summary of this chapter . . . . .	77
<b>6</b>	<b>Discussion</b>	<b>78</b>
6.1	Comparison with theoretical results . . . . .	78
6.1.1	Classical Kitaev model . . . . .	78
6.1.2	Other calculations for the quantum Kitaev model . . . . .	81
6.2	Comparison with experimental results . . . . .	83
6.2.1	Dynamical spin structure factor . . . . .	83
6.2.2	NMR relaxation rate . . . . .	86
6.2.3	Magnetic susceptibility . . . . .	89
6.3	Summary of this chapter . . . . .	90
<b>7</b>	<b>Summary</b>	<b>92</b>
<b>A</b>	<b>Kitaev Hamiltonian in reciprocal space</b>	<b>96</b>
<b>B</b>	<b>Plots of <math>1/T_1</math> and <math>\chi</math> in the <math>T</math>-linear scale</b>	<b>98</b>
<b>C</b>	<b>Benchmark of the maximum entropy method</b>	<b>102</b>
<b>D</b>	<b>Low-<math>T</math> behavior of <math>\chi</math></b>	<b>106</b>
<b>E</b>	<b>Convergence of the hybridization expansion in QMC+CTQMC calculations</b>	<b>108</b>