

Contents

Acknowledgement	i
Abstract	ii
1 Introduction	1
1.1 Multiple Degrees of Freedom in Strongly Correlated Electron Systems	1
1.2 Geometrical Frustration in Lattice Structures	1
1.3 Mixed Valence Spinels	3
1.3.1 AlV ₂ O ₄	4
1.3.2 CuIr ₂ S ₄	5
1.3.3 LiMn ₂ O ₄	6
1.3.4 LiV ₂ O ₄	8
1.4 Purpose of This Thesis	9
1.5 Organization of This Thesis	9
2 Model and Method	11
2.1 Local Density Approximation and Maximally Localized Wannier Function	11
2.2 Multiband Hubbard Model	13
2.2.1 Electron Hopping	14
2.2.2 Trigonal Crystal-field Splitting	14
2.2.3 Relativistic Spin-orbit Interaction	15
2.2.4 Electron-electron Interaction	16
2.3 Generalized Susceptibility	17
2.4 Random Phase Approximation	20
2.5 Fluctuation Analysis	23
3 Result and Discussion	25
3.1 AlV ₂ O ₄	25
3.1.1 Electronic Band Structure	25
3.1.2 Bare Susceptibility	29
3.1.3 RPA Susceptibility	30
3.1.4 Discussion	36
3.2 LiV ₂ O ₄	36
3.2.1 Electronic Band Structure	36
3.2.2 Bare Susceptibility	40
3.2.3 RPA Susceptibility	46
3.2.4 Discussion	51
3.3 Comparison between AlV ₂ O ₄ and LiV ₂ O ₄	53
3.3.1 Electronic Band Structure	53
3.3.2 Tight-binding Parameters and Fermi surfaces	53
3.3.3 Bare Susceptibility	53
3.3.4 Effect of electron correlations: RPA Susceptibility	54
4 Summary	55
A Benchmark for a pyrochlore oxide Cd₂Os₂O₇	57